

Communications

- [16] We found that BRP-H is also a polymerization catalyst for other thiophenes, for example, 3-dodecyl-thiophene.
- [17] *Handbook of Oligo- and Polythiophenes* (Ed.: D. Fichou), Wiley-VCH, Weinheim, 1999.
- [18] D. Dini, F. Decker, F. Andrcani, E. Salatelli, F. Hapiot, *Polymer* 2000, 41, 6473.
- [19] In the measurement setup an argon ion laser beam was focused by a 25 × microscope objective to a 50 μm diameter spot on the sample. The fluorescence emission was imaged by a 50 × objective on a CCD camera.
- [20] No UV/Vis spectrum could be recorded from a single vesicle, nor from a suspension of vesicles, because of scattering problems.
- [21] A 0.5 g L⁻¹ solution of PS-PIAT dissolved in THF was injected into a 30 mg L⁻¹ solution of CAL B enzymes; the final water/THF ratio was 12:1 (v/v). After the mixture was left for two days to equilibrate, it was dialyzed to dispose of all nonincluded enzymes.



Somatostatin Mimics

Design and Synthesis of γ-Dipeptide Derivatives with Submicromolar Affinities for Human Somatostatin Receptors

Dieter Seebach,* Laurent Schaeffer, Meinrad Brenner, and Daniel Hoyer


In a previous paper we have shown that simple *N*-acyl-γ-dipeptide amides that resemble a βII' turn of an α-peptide can be designed to form a turn structure in solution (NMR) and in the solid state (X-ray).^[1,2] To see whether such a turn could also be used to mimic a peptide, the biological activity of which rests upon a turn structure carrying functionalized side chains, we have now synthesized compounds 1a–g (Scheme 1), with the side chain of tryptophan in the γ² position of the first and of lysine in the γ¹ position of the second γ-amino acid, and have tested their affinities for the human somatostatin receptors hssr₁₋₅.^[3–6]

The synthesis of γ-dipeptide derivatives 1 commenced with the *N*-Boc-γ-lactams 2 and 3 (Boc = *tert*-butoxycarbonyl), readily available from the corresponding commercial (*R*)-Ala and (*S*)-Lys acids by known procedures.^[1,7] Ring opening (with the Lys derivative after change of side-chain protection, →4), and esterification with Me₃Si(CH₂)₂OH provided the (*R*)-Boc-γ¹-hhAla and Boc-γ¹-hhLys(Bn₂) esters, which were




Expected conformation of 1

Scheme 1. Structural formulae of the γ-peptides 1. In the expected conformation of 1 the red arrow points to a CH₂ group of the H₂N(CH₂)₄ unit, which is placed inside the shielding cone of the aromatic indole ring. Mes = mesitylenesulfonyl, Bn = benzyl, Nap = naphthyl.



(*R*)-2, R = Me
(*S*)-3, R = (CH₂)₄NH(2-Cl-Z)
(*S*)-4, R = (CH₂)₄NBn₂

5, R = (CH₂)₂SiMe₃
6, R = H



7, R¹ = Boc, R² = Me, X = O(CH₂)₂SiMe₃
8, R¹ = R² = H, X = NHMe
9, R¹ = H, R² = Me, X = NHMe

doubly deprotonated and alkylated with 1-mesitylenesulfonyl-3-bromomethylindole and MeX to give the *unlike* γ²-amino acid derivatives 5 and 7, respectively. The ester group in compound 5 with Trp side chain was cleaved (Bu₄NF, →6), and the lysine-derived esters were converted to the methylamides 8 and 9 without and with 2-methyl substitution, respectively (1. Bu₄NF, 2. MeNH₂, 3. F₃CCO₂H). Coupling of the two γ-amino acid derivatives (6 + 8 and 6 + 9), removal of the Boc groups, and acylation with 2-naphthylacetic acid^[8] (4-methylmorpholine, 1-hydroxy-1*H*-benzotriazole, 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide) produced the side-chain-protected *N*-acyl-dipeptide amides 1a and 1b. Deprotection procedures (MeSO₃H, F₃CCO₂H, and Pd/C, H₂) led to the various partially or fully deprotected γ-dipeptide derivatives 1c–1g. All compounds were purified and fully characterized by elemental analyses, specific optical rotations, circular dichroism (CD), IR, and NMR spectroscopy, and mass spectrometry.

[*] Prof. Dr. D. Seebach, Dr. L. Schaeffer, Dr. M. Brenner, Laboratorium für Organische Chemie, Eidgenössische Technische Hochschule, ETH Hönggerberg, 8093 Zürich (Switzerland). Fax: (+41) 1-632-11-44. E-mail: seebach@org.chem.ethz.ch
Dr. D. Hoyer, Novartis Pharma AG, Nervous System Research, S-386-745, 4002 Basel (Switzerland)

EXHIBIT

C

Figure 1. H-peptides 1c field-shifted

A typical analog: side chain inside the shifts between sections c shown in signals ap; 0.6 ppm, c between the chains, as with a tri compound N-naphthyl exhibit a 10000 deg

↑
H/deg cm²

Figure 2. Nc

Chem.

BEST AVAILABLE COPY




Figure 1. High-field part of the 500 MHz 1H NMR spectra of the γ -dipeptides 1c, 1e, 1f, and 1g in CD_3OD . The red arrows point to high-field-shifted $N(CH_2)_4$ resonances.

A typical feature of the turn structure in somatostatin and its analogues is the juxtaposition of the tryptophan and lysine side chains, which places CH_2 groups of the $H_2N(CH_2)_4$ unit inside the shielding cone of the aromatic indole ring (NMR shifts between $\delta = 0.8$ and 0.3 ppm are observed).^[9] High-field sections of the NMR spectra of four γ -dipeptide amides, shown in Figure 1, in which CH_2 signals appear between $\delta = 0.9$ and 0.6 ppm, confirm the proximity between the corresponding side chains, and are thus compatible with a turn conformation of these compounds. The CD spectra of the *N*-naphthylacetyl dipeptide amides 1 exhibit an intensive negative Cotton effect near 200 nm ($[\theta]$ up to $70000 \text{ deg cm}^2 \text{ dmol}^{-1}$), with a

weaker and broader peak near 220 nm ($[\theta]$ up to $30000 \text{ deg cm}^2 \text{ dmol}^{-1}$) (Figure 2); this CD pattern may be taken as another piece of evidence for the presence of a secondary structure.

Probably the most stringent test of the γ -dipeptide structure is the affinity for somatostatin receptors. Binding affinities for the five cloned human receptors $hsst_{1-5}$, expressed in CCL-39 cell lines, were determined by displacement of [^{125}I]LTT-SRIF₂₈ from these receptor proteins.^[10] While the fully protected γ -dipeptide 1d binds to $hsst_1$ and $hsst_2$ with remarkable K_D values of 0.55 and $1.00 \mu M$, respectively, the partially and the fully deprotected γ -dipeptide derivatives 1f and 1g bind to $hsst_3$ with K_D values of 0.51 and $0.87 \mu M$, respectively (Table 1). Intriguingly, the highest affinities (1d/ $hsst_1$, 1f/ $hsst_3$) are observed when the side chain functional groups (3-indolylmethyl and $(CH_2)_4NH_3^+$) are protected by bulky aromatic moieties (*N*-mesitylenesulfonyl and/or -benzyl)!

The results presented here are confirmative, surprising, and promising; they demonstrate that a 14-amino-acid cyclic disulfide hormone, somatostatin, can be mimicked by a simple, designed, low-molecular-weight, open-chain γ -dipeptide derivative (cf. 1g) that contains only three amide bonds; they suggest that hitherto unknown hydrophobic pockets are present in the receptors ($hsst_1$, $hsst_2$, and $hsst_3$), which supposedly house the turn-bound Trp and Lys side chains (cf. 1c, 1d, 1f); and they promise a potential of γ -peptides for the development of peptidase-resistant^[11] peptidomimetic drugs.

Table 1: pK_D Values for γ -peptides 1b–1g at the five $hsst$ receptors expressed in CCL-39 cells and measured by radioligand binding assays with [^{125}I]LTT-SRIF₂₈ as radioligand.^[10,19]

Receptor	1b	1c	1d	1e	1f	1g	Octreotide ^[a]	SRIF ₁₄ ^[a]
$hsst_1$	5.47	6.06	6.26	5.61	5.98	4.73	6.45	9.08
$hsst_2$	<5	<5	5.17	<5	5.01	2.81	9.11	10.06
$hsst_3$	5.53	5.89	6.00	5.73	5.67	5.42	8.60	9.67
$hsst_4$	4.67	5.74	5.92	5.66	5.79	5.44	5.76	8.39
$hsst_5$	4.49	5.01	5.87	5.14	6.29	6.06	7.31	9.01

[a] Submicromolar affinities are highlighted in red. [b] Sandostatin. [c] Somatostatin.




Figure 2. Nonnormalized CD spectra in MeOH (0.2 mm) of the γ -dipeptide derivatives 1b–1g.

Received: September 25, 2002 [Z50242]

- [1] M. Brenner, D. Seebach, *Helv. Chim. Acta* 2001, 84, 2155–2166.
- [2] For a non-peptidic analogue, see: R. W. Hoffmann, *Angew. Chem.* 2000, 112, 2134–2150; *Angew. Chem. Int. Ed.* 2000, 39, 2054–2070.
- [3] For β -tetrapeptides mimicking somatostatin with nanomolar affinities for $hsst_4$, see: K. Gademann, M. Ernst, D. Hoyer, D. Seebach, *Angew. Chem.* 1999, 111, 1302–1304; *Angew. Chem. Int. Ed.* 1999, 38, 1223–1226; K. Gademann, M. Ernst, D. Seebach, D. Hoyer, *Helv. Chim. Acta* 2000, 83, 16–33; K. Gademann, T. Kimmerlin, D. Hoyer, D. Seebach, *J. Med. Chem.* 2001, 44, 2460–2468; D. Seebach, M. Rueping, P. I. Arvidsson, T. Kimmerlin, P.

BEST AVAILABLE COPY

Communications

Micuch, C. Noti, D. Langenegger, D. Hoyer, *Helv. Chim. Acta* 2001, 84, 3503–3510.

- [4] For a review on somatostatin receptors, see: D. Hoyer, H. Luebbert, C. Bruns, *Naunyn-Schmiedeberg's Arch. Pharmacol.* 1994, 350, 441–453; J. P. Hannon, C. Nunn, B. Stolz, C. Bruns, G. Weckbecker, I. Lewis, T. Tröxler, K. Hurth, D. Hoyer, *J. Mol. Neurosci.* 2002, 18, 15–27. Classification and nomenclature of somatostatin receptors: D. Hoyer, G. I. Bell, M. Berelowitz, J. Epelbaum, W. Feniuk, P. P. Humphrey, A. M. O'Carroll, Y. C. Patel, A. Schombrunn, J. E. Taylor, T. Reisine, *Trends Pharmacol. Sci.* 1995, 16, 86–88.
- [5] For a review on somatostatin analogues, see: A. Janicka, M. Zubrzycka, T. Janicki, *J. Pept. Res.* 2001, 58, 91–107.
- [6] Methyl groups in the γ^1 position of the first and in the γ^2 position of the second γ -amino acid do stabilize the turn (see ref. [1] for more details).
- [7] M. Stureina, P. Majer, E. Majerova, T. A. Guerassina, M. A. Eissanstat, *Tetrahedron* 1997, 53, 12867–12874; S. Hanessian, R. Schaum, *Tetrahedron Lett.* 1997, 38, 163–166; T. Hintermann, K. Gademann, B. Jaun, D. Seebach, *Helv. Chim. Acta* 1998, 81, 983–1002.
- [8] The choice of a 2-naphthylacetyl group at the *N*-terminus was inspired by previous observations: The naphthyl group may be considered to mimic the benzyl side chain of a Phe residue, which precedes the Trp in the peptide chain of somatostatin: S. J. Hocart, R. Jain, W. A. Murphy, J. E. Taylor, B. Morgan, D. H. Coy, *J. Med. Chem.* 1998, 41, 1146–1154; A. J. Souers, A. A. Virgilio, A. Rosenquist, W. Feniuk, J. A. Ellman, *J. Am. Chem. Soc.* 1999, 121, 1817–1825.
- [9] B. H. Arison, R. Hirschmann, D. F. Veber, *Bioorg. Chem.* 1978, 7, 447–451; R. M. Freidinger, D. S. Perlow, W. C. Randall, R. Saperstein, B. H. Arison, D. F. Veber, *Int. J. Pept. Protein Res.* 1984, 23, 142–150; C. Wynants, D. Tourwe, W. Kazmierski, V. I. Hruby, G. Van Binst, *Eur. J. Biochem.* 1989, 185, 371.
- [10] J. P. Hannon, C. Petrucci, D. Fehlmann, C. Viollet, J. Epelbaum, D. Hoyer, *Neuropharmacology* 2002, 42, 396–413. CCL-39 stands for Chinese hamster lung fibroblast cells No. 39. [¹²⁵I]LIT-SRIF₂₈ is a somatostatin dimer containing three mutations (Leu², D-Trp²², ¹²⁵I-Tyr), and SRIF is the acronym for somatostatin release inhibiting factor, 28 refers to the number of amino acids in the physiologically active form of the hormone.
- [11] J. Franckenpohl, P. I. Arvidsson, I. V. Schreiber, D. Seebach, *ChemBioChem* 2001, 2, 445–455.

Controlled Electropolymerization

Liquid-Crystal Templating of Conducting Polymers**

James F. Hulvat and Samuel I. Stupp*

In organic electronics, conducting polymers have a number of advantages over small molecules, particularly because of their stability, mechanical properties, and ease of processing.^[1–4] However, performance of some conducting polymers is limited by their high degree of disorder.^[3,4] Molecular ordering improves carrier mobility in organic field-effect transistors and enhances charge injection in organic light-emitting diodes (OLEDs).^[5] For this reason, vapor-sublimated crystalline films of small molecules are often used,^[6] but alternative strategies to obtain molecular ordering would reduce cost and simplify fabrication of organic electronic devices. One possible way to achieve this is through molecular self-organization. Toward this goal we developed an aqueous, low-temperature technique for preparing conducting polymer films in a self-organized template.

Films of poly(3,4-ethyldioxythiophene) (PEDOT) are commonly used as hole injection layers in OLEDs. PEDOT can be polymerized in organic solvents or in an aqueous suspension with a soluble copolymer or surfactant, leading to amorphous films.^[6] We have studied here the formation of PEDOT films by electropolymerization within a liquid crystalline template.

The well-known hexagonal (H1) lyotropic liquid crystal (LC) consists of cylindrical hydrophobic cores parallel to one another and separated by a hydrophilic continuum (see Figure 1 in the Supporting Information).^[7] LCs have been used by us and others to template inorganic minerals as well as in the formation of mesoporous silica.^[8] With regard to conducting polymers, template chemistry has been used to incorporate chains within the channels of mesoporous silica^[9] or in aqueous channels of an inverse hexagonal LC.^[10] These approaches, however, are limited to soluble, chemically polymerized polymers or to water-soluble monomers such as aniline or pyrrole. Our approach described here is novel in two key respects. First, polymerization occurs in the hydrophobic domain of the LC, allowing use of less polar

[*] Prof. S. I. Stupp, J. F. Hulvat
Department of Materials Science and Engineering
Department of Chemistry
Feinberg School of Medicine
Northwestern University
Evanston, IL 60208-3108 (USA)
Fax: (+1) 847-491-3010
E-mail: s-stupp@northwestern.edu

[**] This work made use of the Electron Probe Instrumentation Center at Northwestern University and was funded by a DoE grant (DE-FG02-00ER45810/A001). J.F.H. is supported by an NDEG fellowship. We thank M. Kern for assistance with XRD and Prof. I. Koltov for helpful discussions.

Supporting information for this article is available on the WWW under <http://www.angewandte.org> or from the author.

monomers such as phenylene,^[11] template n of monom incorporated is avoided electronic phases in water in th

Scheme 1. Aqueous templating

with 3 nm defined by s in the Supp hydrophob: ngly solubl be incorpo suggests th: hydrophob (TEAP) as merization change in 1 and TEAP Differentia EDOT- an isotropizati doped gr We pol: old- or IT(procedure, cooled slow allows the I 500 μ m i 1.3 V versu current den 61 is remc Heating to remaining = Figure 1 function of profilometr limited give isotropic sc (Figure 1b) molecules i Figure 1c s

EST AVAILARIF